# The Beaufort Shelfbreak Jet: Structure, Variability, And Fate

Robert S. Pickart
Physical Oceanography Department. MS #21
Woods Hole Oceanographic Institution, Woods Hole, MA 02543
phone: (508) 289-2858 fax: (508) 457-2181 email: rpickart@whoi.edu

Award Number: N00014-98-1-0046 http://www.whoi.edu/science/PO/people/pickartgroup/index.html

### LONG-TERM GOALS

The long-term goal is to understand the dynamics of exchange between the shelf and open ocean. Regionally, we wish to determine the dominant processes leading to shelf-basin exchange in the western Arctic, and the impact of this on the ventilation of the Arctic halocline.

#### **OBJECTIVES**

The main objectives of this study are (1) To determine the mean transport and cross-stream structure of the Beaufort shelfbreak current; (2) To elucidate the temporal and spatial variability of the jet; and (3) To compare the water mass components of the current to the observed offshore eddies in the western Arctic, to help determine their source.

### **APPROACH**

The approach is to use historical hydrographic and current meter data archived at the National Ocean Data Center, concentrating on the area between Barrow Canyon and 140°W (Figure 1a). It is here




Figure 1a. Study area on the Beaufort shelf and slope. The arrow denotes the flow of Pacific water past Barrow Canyon. The hydrographic sections used in the study are those within the blue box.

maintaining the data needed, and c including suggestions for reducing	nection of information is estimated it, completing and reviewing the collecti this burden, to Washington Headqua uld be aware that notwithstanding an OMB control number.	ion of information. Send comments arters Services, Directorate for Info	regarding this burden estimate or formation Operations and Reports	or any other aspect of the 1215 Jefferson Davis	is collection of information, Highway, Suite 1204, Arlington	
REPORT DATE <b>0 SEP 2001</b> 2. REPORT TYPE				3. DATES COVERED <b>00-00-2001 to 00-00-2001</b>		
4. TITLE AND SUBTITLE				5a. CONTRACT NUMBER		
The Beaufort Shelf	Fate	5b. GRANT NUM	5b. GRANT NUMBER			
				5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)				5d. PROJECT NUMBER		
				5e. TASK NUMBER		
				5f. WORK UNIT NUMBER		
				8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)		
				11. SPONSOR/M NUMBER(S)	ONITOR'S REPORT	
12. DISTRIBUTION/AVAII Approved for publ	LABILITY STATEMENT ic release; distributi	on unlimited				
13. SUPPLEMENTARY NO	OTES					
Regionally, we wish Arctic, and the imp	l is to understand the h to determine the d pact of this on the ve	ominant processes	leading to shelf-ba	-		
15. SUBJECT TERMS						
16. SECURITY CLASSIFIC	ATION OF:		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON	
a. REPORT unclassified	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE unclassified	Same as Report (SAR)	4		

**Report Documentation Page** 

Form Approved OMB No. 0704-0188 that the Pacific water flowing northward from Bering Strait reaches the shelfbreak and turns eastward as a shelfbreak current. There have been 42 hydrographic sections done over the last 50 years (Figure 1a), and a handful of current meter mooring deployments. First, an overall mean section is to be computed, followed by an analysis of the variance about this mean, including a comparison to the observed interior eddy field.

#### WORK COMPLETED

All of the synoptic sections have been extracted from the historical database. Using the International Bathymetric Chart of the Arctic Ocean (IBCAO) data set, an average cross-stream bottom profile was then constructed for the region. Each section was then mapped onto this average bathymetric coordinate system, and objectively interpolated onto a uniform grid. Finally, the interpolated synoptic sections were combined to form mean sections of various hydrographic variables. Standard deviation fields were also produced.

### **RESULTS**

The composite hydrographic sections reveal the unique nature of the Beaufort shelfbreak. In the upper layer (< 50 m) there are two distinct temperature extrema (Figure 1b, color): a narrow core of warm water at the shelfbreak, and a broader area of warmer water further offshore. The inshore signal has a much higher standard deviation, which indicates that warm Bering Strait Summer Water is present intermittently at the shelfbreak. By contrast, the cold subsurface water mass just seaward of the upper slope (centered near 100 m) is more stationary. The origin of this water is still under investigation. The mean potential density section (Figure 1b, contours) reveals a baroclinic jet situated against the upper slope. The sense of the shear implies maximum baroclinic eastward flow near 150-m depth. The current meter data will be used to quantify this.




Figure 1b. Mean section of potential temperature (color), overlaid on potential density (contours).

## **IMPACT/APPLICATIONS**

The mean hydrographic sections have demonstrated the existence of the Beaufort shelfbreak jet, implying that much of the Bering Strait Pacific water "turns the corner" at Barrow Canyon and flows eastward. This suggests that the export directly out of Barrow Canyon into the open Arctic may be less than previously thought. Alternative exchange mechanisms that could transport the water offshore will be investigated in the future using these historical data.

### **TRANSITIONS**

None

## **RELATED PROJECTS**

This project is being carried out in conjunction with the western Arctic Shelf-Basin Interactions Program (jointly sponsored by NSF and ONR).

### **REFERENCES**

None.

# **PUBLICATIONS**

None.